



PROGRAMME



FOR LEAFY VEGETABLE CROPS

Planting	Post-planting	Leaf development	Head formation	Harvest
NHDelta	NHDelta	NHDelta	NHDelta	
XStress	XStress	XStress	XStress	
CalFlux	CalFlux	CalFlux	CalFlux	CalFlux
		RainbowWave		
	TipTop		TipTop	

NHDeltaCa NHDeltaK	After planting, application via irrigation or foliar sprays with NHDeltaCa (5 l/ha) to obtain a better rooting and foliar development decreasing post-transplant stress. From development stage, applications with NHDeltaCa and NHDeltaK to improve leaf growth and quality and a better root system. The use of NHDeltaK improves leaf development for heavier head weight. Nitrate applications can be reduced by up to 20% if required.
XStress	After planting apply at 1.0 to 2.0 l/ha with NHDelta to improve plant health, reduce stress and remove micronutrient deficiencies. From shoot development, regular applications every 10 to 15 days reduce the effects of environmental stress to improve growth and health.
CalFlux	After planting apply at 1.0 l/ha regularly to obtain better developed heads with higher weight, and improved quality; fewer physiological disorders (tip burn, soft tissue, mechanical damage) and increased disease resistance (botrytis, mildew) through improved cell structure. Works synergistically with XStress to reduce stress. Before harvest, 2 applications at 1.5 l/ha; the first 10 days before harvest and the second 3 days to improve post-harvest condition and shelf life.
RainbowWave	Use along growth cycle at 2 l/ha every 12-15 days to decrease vegetative growth.
TipTop	For enhanced growth, increasing both yield and quality. Use as necessary between 2.0 - 3.0 l/ha.