



PROGRAMME FOR ONION CROPS

Planting	Leaf development	Bulb development	Flower bud	Flowering	Bulb development	Harvest
NHDelta	NHDelta	NHDelta	NHDelta	NHDelta	NHDelta	
XStress	XStress	XStress	XStress	XStress	XStress	XStress
					CalFlux	CalFlux
			RainbowWave		RainbowWave	
TipTop		TipTop		TipTop	TipTop	

NHDeltaCa NHDeltaK	After planting, application via irrigation or foliar sprays with NHDeltaCa (5 l/ha) to enhance rooting and foliar development and decreasing post-transplant stress. From development stage, applications with NHDeltaCa and NHDeltaK to improve plant growth and increase bulb weight. Nitrate applications can be reduced by up to 20% if required.
XStress	After planting apply at 1.0 to 2.0 l/ha with NHDelta to improve plant health, reduce stress and remove micronutrient deficiencies. From shoot development, regular applications every 10 to 15 days reduce the effects of environmental stress to improve growth and health.
CalFlux	Use to improve tissue quality and reduce internal disorders caused by calcium deficiency. Works synergistically with XStress to reduce stress and improve plant health. Can be used pre-harvest, 2 applications at 1.5 l/ha; first 10 days and the second 3 days pre-harvest to improve shelf life.
RainbowWave	Use through the growth cycle at 2 l/ha every 12-15 days to decrease vegetative growth and improve bulb and stem weight.
TipTop	For enhanced growth, increasing both yield and quality. Use as necessary between 2.0 - 3.0 l/ha.

Use **TipTop** @ 5.0 l/ha as an alternative to **NHDelta** + **XStress** if plant stress is not an issue. For more information contact your local distributor.